

QUESTION PAPER CODE 65/1/B  
EXPECTED ANSWERS/VALUE POINTS

**SECTION - A**

		Marks
1.	$x = 2, y = 9$ $\therefore x + y = 11$	( $\frac{1}{2}$ for correct x or y) $\frac{1}{2}$ m
2.	order 3, or degree 1 $\therefore$ Degree + order = 4	$\frac{1}{2}$ m $\frac{1}{2}$ m
3.	$\frac{dy}{dx} + \frac{y}{x \log x} = \frac{2}{x}$ (Standard form)  I.F. = $\log x$	$\frac{1}{2}$ m  $\frac{1}{2}$ m
4.	$\vec{a} \cdot \vec{b} = 0 \Rightarrow x = -6$  $y = \pm \sqrt{40}$ or $\pm 2\sqrt{10}$	$\frac{1}{2}$ m  $\frac{1}{2}$ m
5.	$a^2 \sin^2 \alpha + a^2 \sin^2 \beta + a^2 \sin^2 \gamma$ $= 2a^2$	$\frac{1}{2}$ m $\frac{1}{2}$ m
6.	using $\sin \theta = \frac{\vec{a} \cdot \vec{b}}{ \vec{a}  \cdot  \vec{b} }$  $\Rightarrow \theta = 0^\circ$	$\frac{1}{2}$ m  $\frac{1}{2}$ m

**SECTION - B**

7.	$[15000 \quad 15000] \begin{bmatrix} \frac{2}{100} \\ \frac{x}{100} \end{bmatrix} = [1800]$	2 m
----	---	-----

$$\Rightarrow 300 + 150x = 1800 \quad 1 \text{ m}$$

$$\Rightarrow x = 10\%$$

yes : compassionate or any other relevant value 1 m

$$8. \quad \cot^{-1}(x+1) = \sin^{-1} \frac{1}{\sqrt{1+(x+1)^2}} \quad 1\frac{1}{2} \text{ m}$$

$$\text{and } \tan^{-1}x = \cos^{-1} \frac{1}{\sqrt{1+x^2}} \quad 1\frac{1}{2} \text{ m}$$

$$\therefore \sin \left( \sin^{-1} \frac{1}{\sqrt{1+(x+1)^2}} \right) = \cos \left( \cos^{-1} \frac{1}{\sqrt{1+x^2}} \right)$$

$$\Rightarrow 1+x^2+2x+1 = 1+x^2 \Rightarrow x = -\frac{1}{2} \quad 1 \text{ m}$$

OR

$$2 \sin^{-1} \frac{3}{5} - \tan^{-1} \frac{17}{31}$$

$$= 2 \tan^{-1} \frac{3}{4} - \tan^{-1} \frac{17}{31} \quad 1 \text{ m}$$

$$= \tan^{-1} \frac{24}{7} - \tan^{-1} \frac{17}{31} \quad 1 \text{ m}$$

$$= \tan^{-1} 1 = \frac{\pi}{4} \quad 1+1 \text{ m}$$

$$9. \quad C_1 \rightarrow C_1 + C_2 + C_3,$$

$$(a+b+c) \begin{vmatrix} 1 & b & c \\ 1 & c & a \\ 1 & a & b \end{vmatrix} = 0 \quad 1 \text{ m}$$

$$R_2 \rightarrow R_2 - R_1, \quad R_3 \rightarrow R_3 - R_1$$

$$\Rightarrow \begin{vmatrix} 1 & b & c \\ 0 & c-b & a-c \\ 0 & a-b & b-c \end{vmatrix} = 0 \quad (\because a+b+c \neq 0) \quad 2 \text{ m}$$

$$\Rightarrow -a^2 - b^2 - c^2 + ab + bc + ca = 0 \quad \frac{1}{2} \text{ m}$$

$$\Rightarrow -\frac{1}{2} [(a-b)^2 + (b-c)^2 + (c-a)^2] = 0 \quad \frac{1}{2} \text{ m}$$

$$\Rightarrow a = b = c$$

$$10. \quad \begin{pmatrix} 1 & -1 & 0 \\ 2 & 5 & 3 \\ 0 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot A \quad 1 \text{ m}$$

$$R_2 \rightarrow R_2 - 2R_1,$$

$$\begin{pmatrix} 1 & -1 & 0 \\ 0 & 7 & 3 \\ 0 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot A$$

$$R_2 \rightarrow R_2 - 3R_3$$

$$\begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & -3 \\ 0 & 0 & 1 \end{pmatrix} \cdot A$$

$$R_1 \rightarrow R_1 + R_2, \quad R_3 \rightarrow R_3 - 2R_2$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} -1 & 1 & -3 \\ -2 & 1 & -3 \\ 4 & -2 & +7 \end{pmatrix} \cdot A$$

(2 marks for all operations)

$$\therefore A^{-1} = \begin{pmatrix} -1 & 1 & -3 \\ -2 & 1 & -3 \\ 4 & -2 & 7 \end{pmatrix} \quad 1 \text{ m}$$

$$11. \quad f(x) = x - |x - x^2| = |x - x(1-x)| = \begin{cases} 2x - x^2, & -1 \leq x < 0 \\ 0, & x = 0 \\ x^2, & 0 < x \leq 1 \end{cases} \quad 1 \text{ m}$$

$f(x)$  being a polynomial is continuous on  $[-1, 0] \cup [0, 1]$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (2x - x^2) = 0 \quad \frac{1}{2} \text{ m}$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x^2 = 0 \quad \frac{1}{2} \text{ m}$$

$$\text{Also, } f(0) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = f(0) = \lim_{x \rightarrow 0^+} f(x) \quad 1 \text{ m}$$

$\Rightarrow$  There is no point of discontinuity on  $[-1, 1]$  1 m

$$12. \quad \frac{y}{x} = [\log x - \log(a + bx)] \quad \frac{1}{2} \text{ m}$$

$$\Rightarrow \frac{x \frac{dy}{dx} - y}{x^2} = \frac{1}{x} - \frac{b}{a + bx} \quad 1 \text{ m}$$

$$\Rightarrow x \frac{dy}{dx} - y = \frac{ax}{a + bx} \dots\dots\dots (i) \quad 1 \text{ m}$$

Differentiating again,

$$x \frac{d^2y}{dx^2} = \frac{a^2}{(a + bx)^2} \quad 1 \text{ m}$$

$$x^3 \cdot \frac{d^2y}{dx^2} = \left( \frac{ax}{a + bx} \right)^2 = \left( x \frac{dy}{dx} - y \right)^2 \quad (\text{using (i)}) \quad \frac{1}{2} \text{ m}$$

$$13. \quad u = \sec^{-1} \left( \frac{1}{2x^2 - 1} \right) = 2 \cos^{-1} x \Rightarrow \frac{du}{dx} = \frac{-2}{\sqrt{1-x^2}} \quad 1\frac{1}{2} \text{ m}$$

$$v = \sqrt{1-x^2} \Rightarrow \frac{dv}{dx} = \frac{-x}{\sqrt{1-x^2}} \quad 1 \text{ m}$$

$$\left. \frac{dv}{dx} \right|_{x=\frac{1}{2}} = \frac{2}{x} = 4 \quad 1\frac{1}{2} \text{ m}$$

14. Let  $I = \int_0^{\pi/2} \frac{5 \sin x + 3 \cos x}{\sin x + \cos x} dx \dots\dots\dots$  (i)

$\Rightarrow I = \int_0^{\pi/2} \frac{5 \cos x + 3 \sin x}{\cos x + \sin x} dx \dots\dots\dots$  (ii)  $\left( \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right)$  1½ m

Adding (i) and (ii) 1+1 m

$2I = 8 \int_0^{\pi/2} 1 \cdot dx = 4\pi$  ½ m

$\Rightarrow I = 2\pi$

OR

put  $\log x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$  1 m

$= \int e^t \left( \log t + \frac{1}{t^2} \right) dt$

$= \int e^t \left[ \left( \log t - \frac{1}{t} \right) + \left( \frac{1}{t} + \frac{1}{t^2} \right) \right] dt$  1½ m

$= e^t \left( \log t - \frac{1}{t} \right) + c$  1 m

$= x \left[ \log(\log x) - \frac{1}{\log x} \right] + c$  ½ m

15.  $I = \int \frac{x \cos x}{\cos x + x \sin x} dx$  1 m

put  $\cos x + x \sin x = t$

$\Rightarrow x \cos x dx = dt$  1 m

$= \int \frac{dt}{t}$  1 m

$= \log |\cos x + x \sin x| + c$  1 m

$$16. \int \frac{x^4 dx}{(x-1)(x^2+1)} = \int \left[ (x+1) + \frac{1}{(x-1)(x^2+1)} \right] dx \quad 1 \text{ m}$$

(using partial fractions)

$$= \int (x+1) dx + \frac{1}{2} \int \frac{dx}{(x-1)} - \frac{1}{2} \int \frac{x+1}{x^2+1} dx \quad 1\frac{1}{2} \text{ m}$$

$$= \frac{x^2}{2} + x + \frac{1}{2} \log |x-1| - \frac{1}{4} \log (x^2+1) - \frac{1}{2} \tan^{-1} x + c \quad 1\frac{1}{2} \text{ m}$$

$$17. \left. \begin{aligned} \overline{AB} &= -2\hat{i} - 5\hat{k} \\ \overline{AC} &= \hat{i} - 2\hat{j} - \hat{k} \end{aligned} \right\} \quad 1 \text{ m}$$

$$\overline{AB} \times \overline{AC} = -10\hat{i} - 7\hat{j} + 4\hat{k} \quad 1 \text{ m}$$

$$|\overline{AB} \times \overline{AC}| = \sqrt{165} \quad \frac{1}{2} \text{ m}$$

$$\hat{n} = \frac{\overline{AB} \times \overline{AC}}{|\overline{AB} \times \overline{AC}|} \quad 1 \text{ m}$$

$$= \frac{(-10\hat{i} - 7\hat{j} + 4\hat{k})}{\sqrt{165}} \text{ or } \frac{10\hat{i} + 7\hat{j} - 4\hat{k}}{\sqrt{165}} \quad \frac{1}{2} \text{ m}$$

$$18. \left. \begin{aligned} \vec{a}_1 &= -\hat{i}, \vec{b}_1 = \hat{i} + \frac{1}{2}\hat{j} - \frac{1}{12}\hat{k} \\ \vec{a}_2 &= -2\hat{j} + \hat{k}, \vec{b}_2 = \hat{i} + \hat{j} + \frac{1}{6}\hat{k} \end{aligned} \right\} \quad 1 \text{ m}$$

$$\vec{a}_2 - \vec{a}_1 = \hat{i} - 2\hat{j} + \hat{k} \quad \frac{1}{2} \text{ m}$$

$$\vec{b}_1 \times \vec{b}_2 = \frac{1}{6}\hat{i} - \frac{1}{4}\hat{j} + \frac{1}{2}\hat{k} \quad \frac{1}{2} \text{ m}$$

$$|\vec{b}_1 \times \vec{b}_2| = \frac{7}{12} \quad 1 \text{ m}$$

$$\text{S.D.} = \left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right| = 2 \quad 1 \text{ m}$$

OR

Foot of perpendicular are  $(0, b, c)$  &  $(a, 0, c)$  1 m  
Equ. of required plane

$$\begin{vmatrix} x & y & z \\ 0 & b & c \\ a & 0 & c \end{vmatrix} = 0 \quad 2 \text{ m}$$

$$\Rightarrow bcx + acy - abz = 0 \quad 1 \text{ m}$$

19.  $p(x=2) = 9 \cdot P(x=3)$  1 m

$$\Rightarrow {}^3C_2 p^2 q = 9 \cdot {}^3C_3 p^3 \cdot q^0 \quad 1 \text{ m}$$

$$\Rightarrow 3p^2(1-p) = 9p^3 \quad 1 \text{ m}$$

$$\Rightarrow p = \frac{1}{4} \quad 1 \text{ m}$$

OR

Let  $H_1$  be the event that red ball is drawn

$H_2$  be the event that black ball is drawn

E be the event that both balls are red

$$P(H_1) = \frac{3}{8}, \quad P(H_2) = \frac{5}{8} \quad 1 \text{ m}$$

$$P(E/H_1) = \frac{{}^5C_2}{{}^{10}C_2} = \frac{2}{9}, \quad P(E/H_2) = \frac{{}^3C_2}{{}^{10}C_2} = \frac{1}{15} \quad 1 \text{ m}$$

$$P(E) = P(H_1) P(E/H_1) + P(H_2) \cdot P(E/H_2) \quad 1 \text{ m}$$

$$= \frac{3}{8} \cdot \frac{2}{9} + \frac{5}{8} \cdot \frac{1}{15} = \frac{1}{8} \quad 1 \text{ m}$$

20.

*	0	1	2	3	4	5	6
0	0	1	2	3	4	5	6
1	1	2	3	4	5	6	0
2	2	3	4	5	6	0	1
3	3	4	5	6	0	1	2
4	4	5	6	0	1	2	3
5	5	6	0	1	2	3	4
6	6	0	1	2	3	4	5

4 m

$$\forall a \in \{0, 1, 2, 3, 4, 5, 6\}$$

$$a * 0 = a = 0 * a \Rightarrow 0 \text{ is identity}$$

1 m

$$\forall a \in \{1, 2, 3, 4, 5, 6\}$$


$$a * b = 0 = b * a$$

$$\Rightarrow a * (7 - a) = 0 = (7 - a) * a$$

$$\Rightarrow (7 - a) \text{ is inverse of } a$$

1 m

21.  $A = y(x + 4)$



$$\frac{x^2}{16} + \frac{y^2}{9} = 1$$

$$\text{Let } z = A^2 = \frac{9}{16}(16 - x^2)(x + 4)^2 \Rightarrow y^2 = \frac{9}{16}(16 - x^2) \dots\dots\dots (i)$$

1 m

$$= \frac{9}{16}(4 - x)(4 + x)^3$$

1 m

$$\frac{dz}{dx} = \frac{9}{16} (4+x)^2 (8-4x) \quad 1 \text{ m}$$


$$\frac{dz}{dx} = 0 \Rightarrow x = 2 \quad 1 \text{ m}$$

$$\frac{d^2z}{dx^2} = -\frac{9}{4} (4+x)^2 + \frac{9}{8} (4+x) (8-4x)$$

$$\left. \frac{d^2z}{dx^2} \right|_{x=2} < 0 \quad 1 \text{ m}$$

$$\therefore \text{Maximum value of } A = 9\sqrt{3} \text{ sq. units} \quad 1 \text{ m}$$

22.



Required Area

$$= \int_0^2 \sqrt{4-x^2} dx - \int_0^2 (2-x) dx \quad 2 \text{ m}$$

$$= \left[ \frac{x\sqrt{4-x^2}}{2} + \frac{4}{2} \sin^{-1} \frac{x}{2} \right]_0^2 - \left[ 2x - \frac{x^2}{2} \right]_0^2 \quad 1+1 \text{ m}$$

$$= (\pi - 2) \text{ sq. units} \quad 1 \text{ m}$$

$$23. \quad \frac{dy}{dx} = \frac{xy}{x^2 + y^2}$$

$$\text{put } y = v x \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad 1 \text{ m}$$

$$\Rightarrow \frac{1+v^2}{v^3} = -\frac{dx}{x} \quad 1 \text{ m}$$

Integrating both sides

$$-\frac{1}{2v^2} + \log v = -\log x + c \quad 1 \text{ m}$$

$$\Rightarrow -\frac{x^2}{2y^2} + \log y = c$$

$$\text{when } x = 1, y = 1 \Rightarrow c = -\frac{1}{2} \quad 1 \text{ m}$$

$$\Rightarrow \log y = \frac{x^2 - y^2}{2y^2} \quad \frac{1}{2} \text{ m}$$

$$\text{when } x = x_0, y = e \Rightarrow x_0 = \sqrt{3} e \quad 1\frac{1}{2} \text{ m}$$

OR

$$I F = e^{\int \tan x \, dx} = e^{\log \sec x} = \sec x \quad 1 \text{ m}$$

$$\therefore \frac{d}{dx} (y \cdot \sec x) = 3x^2 \sec x + x^3 \sec x \tan x \quad 1 \text{ m}$$

$$\Rightarrow y \sec x = \int 3x^2 \sec x \cdot dx + x^3 \sec x - \int 3x^2 \cdot \sec x \, dx + c \quad 2 \text{ m}$$

$$\Rightarrow y = x^3 + c \cos x$$

$$\text{when } x = \frac{\pi}{3}, y = 0; \text{ we get } c = \frac{-2\pi^3}{27} \quad 1 \text{ m}$$

$$\therefore y = x^3 - \frac{2\pi^3}{27} \cos x \quad 1 \text{ m}$$

24. Equation of line is  $\frac{x-3}{2} = \frac{y-4}{-3} = \frac{z-1}{5}$  1 m

Equation of plane is

$$\begin{vmatrix} x-2 & y-1 & z-2 \\ 1 & 0 & -2 \\ 2 & -3 & -1 \end{vmatrix} = 0$$
 1 m

$\Rightarrow 2x + y + z - 7 = 0$  ..... (i) 1 m

general point on given line  $(2\lambda + 3, -3\lambda + 4, 5\lambda + 1)$  lies on (i) 1 m

$\therefore 2(2\lambda + 3) + (-3\lambda + 4) + (5\lambda + 1) - 7 = 0 \Rightarrow \lambda = -\frac{2}{3}$  1 m

$\therefore$  Point of intersection  $\left(\frac{5}{3}, 6, -\frac{7}{3}\right)$  1 m

25. Let  $H_1$  : be the event 1, 2 appears

$H_2$  : be the event 3, 4, 5, 6 appears 1 m

$E_3$  : be the event that head appears

$P(H_1) = \frac{2}{6} = \frac{1}{3}, P(H_2) = \frac{4}{6} = \frac{2}{3}$  1 m

$P(E/H_1) = \frac{3}{8}, P(E/H_2) = \frac{1}{2}$  1 m

$P(H_2/E) = \frac{P(H_2) \cdot P(E/H_2)}{P(H_1) \cdot P(E/H_1) + P(H_2) \cdot P(E/H_2)}$  1 m

$= \frac{8}{11}$  2 m

OR

Let  $H_1$  : be the event that 4 occurs

$H_2$  : be the event that 4 does not occurs 1 m

$E$  : be the event that man reports 4 occurs

on a throw of dice

$$P(H_1) = \frac{1}{6}, \quad P(H_2) = \frac{5}{6} \quad 1 \text{ m}$$

$$P(E/H_1) = \frac{3}{5} \quad P(E/H_2) = 1 - \frac{3}{5} = \frac{2}{5} \quad 1 \text{ m}$$

$$P(H_1/E) = \frac{P(H_1) \cdot P(E/H_1)}{P(H_1) \cdot P(E/H_1) + P(H_2) \cdot P(E/H_2)} \quad 1 \text{ m}$$

$$= \frac{3}{13} \quad 2 \text{ m}$$

26. Let us consider the man invested on x electronic and y manually operated machines


$$\text{Maximise } P = 220x + 180y \dots\dots\dots (i) \quad 1 \text{ m}$$

subject to

$$x + y \leq 20$$

$$3600x + 2400y \leq 57600 \Rightarrow 3x + 2y \leq 48 \quad 1\frac{1}{2} \text{ m}$$

$$x, y \geq 0$$



(1 mark for plotting each line) = 2 m

(½ to find the vertices of feasible region)

$$P \Big|_{A(16,0)} = 3520 \text{ Rs.}$$

$$P \Big|_{B(8,12)} = 3920 \text{ Rs.}$$

$$P \Big|_{C(0,20)} = 3600 \text{ Rs.}$$

Maximum profit is Rs. 3920 at x = 8, y = 12 1 m