






CHEMISTRY MARKING SCHEME

FOREIGN-2016






SET -56/2/3/F

Q.no.	Answers	Marks
1	NO ₂ gas	1
2	N,N-dimethylbutanamide	1
3	Like Charged particles cause repulsion/ Brownian motion/ solvation	1
4	Because of some crystallization.	1
5	Reaction (ii)	1
6	X = CH ₃ -CO-CH ₂ -CH ₃ / Butan-2-one Y = CH ₃ -CH(OH)-CH ₂ -CH ₃ / Butan-2-ol	1 1
7	<p>i) </p> <p>ii) </p>	1+1
8	<p>i) [Co(NH₃)₄Cl₂]Cl</p> <p>ii) Tetraamminedichloridocobalt(III) chloride</p>	1 1
9	<p>When reaction is completed 99.9%, [R]_t = [R]₀ - 0.999[R]₀</p> $k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$ $= \frac{2.303}{t} \log \frac{[R]_0}{[R]_0 - 0.999[R]_0} = \frac{2.303}{t} \log 10^3$ $t = 6.909/k$ <p>For half-life of the reaction</p> $t_{1/2} = 0.693/k$ $\frac{t}{t_{1/2}} = \frac{6.909}{k} \times \frac{k}{0.693} = 10$	1/2 1/2 1
OR		
9	<p>R → P</p> $\text{Rate} = \frac{dR}{dt} = kR$ <p>or $\frac{dR}{R} = -kdt$</p> <p>Integrating this equation, we get</p> $\ln [R] = -kt + I \tag{4.8}$ <p>Again, I is the constant of integration and its value can be determined easily.</p> <p>When t = 0, R = [R]₀, where [R]₀ is the initial concentration of the reactant.</p> <p>Therefore, equation (4.8) can be written as</p> $\ln [R]_0 = -k \times 0 + I$ $\ln [R]_0 = I$ <p>Substituting the value of I in equation (4.8)</p> $\ln [R] = -kt + \ln [R]_0 \tag{4.9}$ <p>Rearranging this equation</p> $\ln \frac{R}{R_0} = -kt$ <p>or $k = \frac{1}{t} \ln \frac{[R]_0}{[R]}$, $k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$</p>	1/2 1/2 1

10	Henry's law states that the mole fraction of gas in the solution is proportional to the partial pressure of the gas over the solution. Applications: solubility of CO ₂ gas in soft drinks /solubility of air diluted with helium in blood used by sea divers or any other Solubility of gas in liquid decreases with increase in temperature.	1 1/2 1/2
11	(i) Butadiene and acrylonitrile CH₂ = CH – CH = CH₂ and CH₂=CH-CN (ii) Vinyl chloride CH₂=CH-Cl (iii) Chloroprene Cl CH₂ = C – CH = CH₂	1/2+1/2 1/2+1/2 1/2+1/2
12	i)  ii) Peptide linkage / -CO-NH- linkage iii) Water soluble-Vitamin B / C Fat soluble- Vitamin A /D /E /K	1 1 1/2+1/2
13	i) dsp ³ , Diamagnetic, low spin ii) The energy used to split degenerate d-orbitals due to the presence of ligands in a definite geometry is called crystal field splitting energy.	1 1/2+1/2 1
14	i) Iodine is heated with Zr or Ti to form a volatile compound which on further heating decompose to give pure Zr or Ti . or $\text{Zr(impure)} + 2\text{I}_2 \longrightarrow \text{ZrI}_4 \text{ (volatile)}$ $\text{ZrI}_4 \xrightarrow{1800\text{K}} \text{Zr(pure)} + 2\text{I}_2$ ii) Cryolite lowers the m.p. of alumina mix / acts as a solvent / brings conductivity. iii) Role of NaCN in the extraction of Ag is to do the leaching of silver ore in the presence of air. or $4\text{Ag(s)} + 8\text{CN}^-(\text{aq}) + 2\text{H}_2\text{O} + \text{O}_2(\text{g}) \longrightarrow 4[\text{Ag}(\text{CN})_2]^- + 4\text{OH}^-$	1 1 1 1

15	<p>i)</p>  <p>ii)</p>  <p>iii) $\text{CH}_3\text{CH}_2\text{ONO}$</p>	1 x 3=3
16	$k = \frac{2.303}{t} \log \frac{p_i}{2p_i - p_t}$ $= \frac{2.303}{300} \log \frac{0.3}{2 \times 0.3 - 0.5}$ $= \frac{2.303}{300} \log 3$ $= \frac{2.303 \times 0.4771}{300}$ $= 0.0036 \text{ atm}^{-1} \text{ or } 0.004 \text{ atm}^{-1} \text{ (approx.)}$	1 1 1
17	<p>i) Because of the resonance stabilization of the conjugate base i.e enolate anion or diagrammatic representation.</p> <p>iii) Because the carboxyl group gets bonded to the catalyst anhyd. AlCl_3 (Lewis acid). (note: part ii is deleted because of printing error and mark allotted in part i and part iii)</p>	1½ 1½
OR		
17	<p>i) $\text{C}_6\text{H}_5\text{CH}_3 \xrightarrow{\text{CrO}_3/(\text{CH}_3\text{CO})_2\text{O}} \text{C}_6\text{H}_5\text{CH}(\text{OCOCH}_3)_2 \xrightarrow{\text{H}_2\text{O}} \text{C}_6\text{H}_5\text{CHO}$</p> <p>ii) $\text{CH}_3\text{COOH} \xrightarrow{\text{Cl}_2/\text{P}} \text{Cl-CH}_2\text{-COOH}$</p> <p>iii) $\text{CH}_3\text{COCH}_3 \xrightarrow{\text{Zn(Hg)/conc.HCl}} \text{CH}_3\text{CH}_2\text{CH}_3$</p> <p style="text-align: center;">(Or by any other correct method)</p>	1x3=3

18	$d = \frac{z \times M}{N_A \times a^3}$ <p style="text-align: center;">Or</p> $d = \frac{z \times w}{N \times a^3} \quad \text{Where } w \text{ is weight and } N \text{ is no. of atoms.}$ $d = \frac{4 \times 200 \text{ g}}{2.5 \times 10^{24} \times (400 \times 10^{-10} \text{ cm})^3}$ $d = 5 \text{ g cm}^{-3}$ <p style="text-align: center;">(or by any other correct method)</p>	1 1 1
19	<p>i) It is a process in which both adsorption and absorption can take place simultaneously.</p> <p>ii) It is the potential difference between the fixed layer and the diffused/ double layer of opposite charges around the colloidal particles.</p> <p>iii) It is the temperature above which the formation of micelles takes place.</p>	1 1 1
20	$\Delta T_f = iK_f m$ <p>For complete ionisation of Na_2SO_4 $i=3$</p> $\Delta T_f = T_f^0 - T_f = 3 \times 1.86 \text{ K kg mol}^{-1} \times \frac{2\text{g}}{142\text{g mol}^{-1}} \times \frac{1000 \text{ g kg}^{-1}}{50 \text{ g}}$ $\Delta T_f = 1.57$ <p style="text-align: center;">So, $T_f = -1.57^\circ\text{C}$ or 271.43K</p>	$\frac{1}{2}$ $\frac{1}{2}$ 1 1
21	<p>i) Because of higher oxidation state (+5) / high charge to size ratio / high polarizing power.</p> <p>ii) Because of high interelectronic repulsion.</p> <p>iii) Because of its low bond dissociation enthalpy and high hydration enthalpy of F^-.</p>	$1 \times 3 = 3$
22	<p>i) A : $\text{C}_6\text{H}_5\text{CONH}_2$ B : $\text{C}_6\text{H}_5\text{NH}_2$ C : $\text{C}_6\text{H}_5\text{NHCOCH}_3$</p> <p>ii) A: $\text{C}_6\text{H}_5\text{NO}_2$ B : $\text{C}_6\text{H}_5\text{NH}_2$ C: $\text{C}_6\text{H}_5\text{-NC}$</p>	$1\frac{1}{2}$ $1\frac{1}{2}$
23	<p>(i) Caring ,dutiful, Concerned, compassionate (or any other two values)</p> <p>ii) Because higher doses may have harmful effects and act as poison which cause even death.</p> <p>iii) Tranquilizers are a class of chemical compounds used for treatment of stress or even mental diseases. ex. chlordiazepoxide, equanil, veronal, serotonin, valium (or any other two examples)</p>	$\frac{1}{2} + \frac{1}{2}$ 1 1 $\frac{1}{2} + \frac{1}{2}$

24	<p>a)</p> <p>i) $(\text{CH}_3)_3\text{C-I} + \text{CH}_3\text{-OH}$</p> <p>ii) $\text{CH}_3\text{-CH}_2\text{-C(=O)-CH}_3$</p> <p>iii)</p>  <p>b) i)</p>  <p>ii).</p> 	1 1 1 1
OR		
24	<p>a).</p> <p>(i)</p>  <p>(ii) $\text{CH}_3\text{CH}_2\text{OH} + \text{CH}_3\text{COCl} \xrightarrow{\text{pyridine}} \text{CH}_3\text{CH}_2\text{O-COCH}_3 + \text{HCl}$</p> <p>(iii).</p> 	1 1 1

	(b)(i) Warm each compound with iodine and sodium hydroxide. Phenol : No yellow ppt formed Ethanol: Yellow ppt of Iodoform are formed.	1
	ii) On adding Lucas reagent (HCl/anhyd. ZnCl ₂), Propan-2-ol gives white turbidity after 5 minutes whereas 2-methylpropan-2-ol gives white turbidity immediately. (or any other suitable test)	1

25	a) Given $E^{\circ}_{\text{Cell}} = + 0.30\text{V}$; $F = 96500\text{C mol}^{-1}$ $n = 6$ (from the given reaction) $\Delta_r G^{\circ} = - n \times F \times E^{\circ}_{\text{Cell}}$ $\Delta_r G^{\circ} = - 6 \times 96500 \text{ C mol}^{-1} \times 0.30\text{V}$ $= - 173,700 \text{ J / mol or } - 173.7 \text{ kJ / mol}$ $\log Kc = \frac{n E^{\circ}_{\text{Cell}}}{0.059}$ $\log Kc = \frac{6 \times 0.30}{0.059}$ $\log Kc = 30.5$	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1
	b) A Because E° value of A shows that on coating ,A acts as anode and Fe acts as a cathode and hence A oxidises in preference to Fe and prevent corrosion / or E°_{cell} is positive and hence A oxidises itself to prevent corrosion of Fe/ E° value is more negative. (or any other correct reason)	1 1 1

OR

25	a) $\Lambda_m = \frac{\kappa}{c}$ $= \frac{3.905 \times 10^{-5} \text{ S cm}^{-1}}{0.001 \text{ mol L}^{-1}} \times \frac{1000 \text{ cm}^3}{\text{L}}$ $\Lambda_m = 39.05 \text{ Scm}^2 \text{ mol}^{-1}$ $\Lambda_o = \lambda^{\circ}(\text{H}^+) + \lambda^{\circ}(\text{CH}_3\text{COO}^-)$ $= (349.6 + 40.9) \text{ Scm}^2 \text{ mol}^{-1}$ $\Lambda_o = 390.5 \text{ Scm}^2 \text{ mol}^{-1}$ $\alpha = \frac{\Lambda_m}{\Lambda_o}$ $= \frac{39.05 \text{ Scm}^2 \text{ mol}^{-1}}{390.5 \text{ Scm}^2 \text{ mol}^{-1}}$ $\alpha = 0.1$	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1
----	--	--

	b)Secondary battery or rechargeable battery	1
	$\text{Pb(s)} + \text{PbO}_2\text{(s)} + 2\text{SO}_4^{2-}\text{(aq)} + 4\text{H}^+\text{(aq)} \longrightarrow$ $2\text{PbSO}_4\text{(s)} + 2\text{H}_2\text{O(l)}$	1

26	a) <ul style="list-style-type: none"> i)Because of higher oxidation state (+7) of Mn. 1 ii)Because it has one unpaired electron in 3d orbital in its +2 oxidation state / or it has incompletely filled d-orbital in +2 oxidation state. 1 iii)Because of comparable energies of 5f , 6d and 7s orbitals. 1 b) $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \longrightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$ $3\text{MnO}_4^{2-} + 4\text{H}^+ \longrightarrow 2\text{MnO}_4^- + \text{MnO}_2 + 2\text{H}_2\text{O}$	1+1
----	--	-----

OR

26	a) <ul style="list-style-type: none"> i)Cr, because of maximum no. of unpaired electrons cause strong metallic bonding. $\frac{1}{2} + \frac{1}{2}$ ii)Mn, because it attains stable half -filled 3d⁵ configuration in +2 oxidation state. $\frac{1}{2} + \frac{1}{2}$ iii)Zn, because of no unpaired electron in d-orbital. $\frac{1}{2} + \frac{1}{2}$ b) $2\text{Na}_2\text{CrO}_4 + 2\text{H}^+ \rightarrow \text{Na}_2\text{Cr}_2\text{O}_7 + 2\text{Na}^+ + \text{H}_2\text{O}$ $\text{Na}_2\text{Cr}_2\text{O}_7 + 2\text{KCl} \longrightarrow \text{K}_2\text{Cr}_2\text{O}_7 + 2\text{NaCl}$	1+1
----	--	-----